

The **Irish** Longitudinal Study on Ageing

Derived Variables Codebook

Wave 5

Irish Longitudinal Study on Ageing,
Trinity Central,
152-160 Pearse St,
Dublin 2,
D02 R590,
Republic of Ireland

Telephone: +353 1 896 2509 Fax: +353 1 896 2451

The Irish Longitudinal Study on Ageing is funded by:

This codebook contains information on additional variables included in the TILDA Wave 5 dataset. Notes on how the variable was derived are included where necessary.

A list of the original variables used for derivation are included. Please note that not all original variables are included in released versions of the datasets.

Where possible, details of any original documentation relating to scale scoring or information are included.

For any further enquires on variables not listed here, please refer to the TILDA release guide available on the TILDA website www.tilda.ie.

<http://www.tilda.ie>



The Irish Longitudinal
Study on Ageing

Contents

Functional Limitations	8
ADL_reported	8
ADL_helper	8
IADL_reported.....	8
IADL_helper	9
adla.....	9
adlwa	11
iadla.....	11
iadlza.....	12
mobilb	12
lgmusa.....	12
grossa	13
finea	13
DISimpairments	13
DISadl	13
DISdisab3.....	14
DISdisab4.....	14
DISlongterm.....	14
DISvision	14
DIShearing_withaid	15
DIShearing_withoutaid	15
DISconverse4.....	16
DIStaste.....	16
DISsmell	16
Health.....	17
COGimmediatedrecall1	17
COGimmediatedrecall2	17
COGdelayedrecall.....	17
COGprosmem1	17
COGprosmem2	18
COGmmse	18
COGanimal_naming	18
MHcesd_capi_sf	19
MHpennworry.....	19
MHucla_loneliness	20
MHcapi_loneliness	22

MHcidi_anxiety	22
MHcidi_depression	23
MHcasp12_total	24
MHcasp12_control	25
MHcasp12_autonomy	25
MHcasp12_pleasure	26
MHcasp12_selfreal	26
PSSscore	27
CHRany_pain	28
CHRpain_severity	28
CHRpainlimits	28
MDmeds	29
MDmeds4	29
MDmeds_excl_supps	29
MDmeds4_excl_supps	29
MDsupplements	30
MDpolypharmacy	30
MDpoly_excl_supps	30
MDpoly_extreme	31
MDpoly_extreme_ex_supps	31
MDantidepressant	31
MDantihypertensives	32
MDantihypertensives2	32
medcard	32
MDace_bbb	33
MDace_nonbbb	33
MDarb_bbb	33
MDarb_nonbbb	33
MDarb_undbbb	34
MDccb_bbb	34
MDccb_nonbbb	34
MDantihypertensivesC08CEG	35
MDhypoglyc	35
MDinsulin	35
MDglycosides	36
Mdantiarrhythmics	36
MDvasodil	36

MDantiadren	37
MDdiur.....	37
MDlc_diur.....	37
MDhc_diur.....	38
MDnons_betab.....	38
MDs_betab.....	38
MDccb	39
MDace.....	39
MDarb.....	39
MDstatin	40
MDincont	40
MDalpha_bph	40
MDnsaid.....	41
MDopiod.....	41
MDantiepileptic	41
MDantipark	Error! Bookmark not defined.
MDantipsych.....	42
MDnons_mari	42
MDssri	42
MDothertidep	43
MDantihist	43
MDanticholinergic	43
smokcurr	44
BEHcage	45
BEHcage2	45
ICD Codes.....	46
ICD10_01.....	46
ICD10_02.....	46
ICD10_03.....	47
ICD10_04.....	47
ICD10_05.....	47
ICD10_06.....	48
ICD10_07.....	48
ICD10_08.....	48
ICD10_09.....	49
ICD10_10.....	49
ICD10_11.....	49

ICD10_12.....	50
ICD10_13.....	50
ICD10_14.....	50
ICD10_15.....	51
ICD10_16.....	51
IPAQ	52
minswalking.....	52
minsmoderate.....	52
minsvigorous.....	52
walkingmet.....	53
moderatemet.....	53
vigorousmet.....	53
FRipaquetminutes	53
FRexercise3.....	54
Employment.....	55
retired	55
employ	55
pension_payIn.....	55
pension_receive.....	56
Financial.....	57
INCASSassets.....	57
INCASSnetassets	57
INCASSassets5	57
INCASSnetassets5	57
INCtotalhousehold_loop	57
INCtotalhousehold_loop	58
Social Participation	59
SOCpaliveW5	59
SOCsocParticip.....	59
SOCChurch.....	59
SOCClubs	60
SOCrelFriends.....	60
SOCcloseTies.....	60
SOCsocnet	60
SCR Section.....	61
SCRflu	61
SCRbreastlumps.....	61

Functional Limitations

Activities of Daily Living

ADL_reported

Description: Indicator variable for respondents who have reported any ADL.

Notes: *This variable is not comparable with TILDA Wave 1.*

Response options:

0 = No

1 = Yes

98 = Don't Know

99 = Refused

Variables used:

f1002_01 - f1002_06

ADL_helper

Description: Indicator variable for respondents who report having help for their ADL(s).

Notes: *This variable is not comparable with TILDA Wave 1.*

Response options:

0 = No

1 = Yes

98 = Don't Know

99 = Refused

Variables used:

f1005, f1008, f1011, f1013, f1016, f1019

IADL_reported

Description: Indicator variable for respondents who have reported any IADLs.

Response options:

0 = No

1 = Yes

98 = Don't Know

99 = Refused

Variables used:

f1025_01 - f1025_06

IADL_helper

Description: Indicator variable for respondents who report having help for their IADL(s).

Response options:

0 = No

1 = Yes

98 = Don't Know

99 = Refused

Variables used:

f1026, f1029, f1030, f1031

adla

Description: Reports number of ADLs from list of 5.

ADLs included:

- Difficulty bathing or showering
- Dressing
- Eating, such as cutting up your food
- Getting in or out of bed
- Walking across a room.

0 = None

1 = Single ADL

2 = 2 ADLs

3 = 3 ADLs

Variables used:

f1002_01 - f1002_05

adlwa

Description: Number of Wallace and Herzog proposed ADLs: 0-3.

ADLs included:

- Difficulty bathing or showering
- Dressing
- Eating, such as cutting up your food

0 = None

1 = Single ADL

2 = 2 ADLs

3= 3 ADLs

Notes:

Refer to Wallace, R. B., & Herzog, A. R. (1995). Overview of the health measures in the Health and Retirement Study. *The Journal of Human Resources* XXX (Suppl.), S84-S107 for more details on the Wallace and Herzog proposed ADL scale.

Variables used:

f1002_01, f1002_03, f1002_04

iadla

Description: Reports number of IADLs from list of 3.

IADLs included:

- Making telephone calls
- Taking medications
- Managing money, such as paying bills and keeping track of expenses

Variables used:

f1025_04 – f1025_06

iadlza

Description: Reports number of IADLs from list of 5.

IADLs included:

- Preparing a hot meal
- Shopping for groceries
- Making telephone calls
- Taking medications
- Managing money, such as paying bills and keeping track of expenses

Variables used:

f1025_01, f1025_03, f1025_04, f1025_05, f1025_06

mobilb

Description: ADL & IADL mobility index: 0-4.

Limitations included:

- Walking 100 meters (100 yards)
- Walking across a room
- Climbing several flights of stairs
- Climbing one flight of stairs without resting

Variables used:

f1001_01, f1001_05, f1001_06, f1002_02

lgmusa

Description: ADL & IADL large muscle index: 0-4.

Limitations included:

- Sitting for about two hours
- Getting up from a chair after sitting for long periods
- Stooping, kneeling or crouching
- Pulling or pushing large objects like a living room chair

Variables used:

f1001_03, f1001_04, f1001_07, f1001_09

grossa

Description: ADL & IADL gross motor index: 0-5.

Limitations included:

- Walking 100 meters
- Walking across a room
- Climbing one flight of stairs without resting for long periods
- Getting in or out of bed
- Bathing or showering

Variables used:

f1001_01, f1001_06, f1002_02, f1002_05, f1002_03

finea

Description: ADL & IADL fine motor: 0-3.

Limitations included:

- Picking up a small coin from a table
- Eating, such as cutting up your food
- Dressing

Variables used:

f1001_11, f1002_01, f1002_04

DISimpairmentsDescription: Number of physical limitations

Variables used:

f1001_01 – f1001_11

DISadl

Description: Number of ADL impairments

Variables Used:

f1002_01 - f1002_06

DISiadl

Description: Number of IADL impairments

Variables Used:

f1025_04 – f1025_06

DISdisab3

Description: Indicator variable for respondents with ADL or IADL disabilities

Variables Used:

f1001_01 – f1001_11, f1002_01 - f1002_06, f1025_01 - f1025_06

DISdisab4

Description: Indicator variable for respondents with ADL or IADL disabilities

Variables Used:

f1001_01 – f1001_11, f1002_01 - f1002_06, f1025_01 - f1025_06

DISlongterm

Description: Indicator for respondents with long term illness

Response Options

1 = No longer term illness

2 = Long term illness

3 = Limiting long term illness

Variables Used:

ph003

DISvision

Description: Respondent's self-rated vision

Response Options:

1 = Excellent

2 = Very Good

3 = Good

4 = Fair

5 = Poor or Registered Blind

Variables Used:

ph102

DIShearing_withaid

Description: Respondent's self-rated hearing with aid

Response Options:

1 = Excellent

2 = Very Good

3 = Good

4 = Fair

5 = Poor or Registered Blind

Variables Used:

ph108b

DIShearing_withoutaid

Description: Respondent's self-rated hearing without aid

Response Options:

1 = Excellent

2 = Very Good

3 = Good

4 = Fair

5 = Poor or Registered Blind

Variables Used:

ph108a

DISconverse4

Description: Difficulty following a conversation with 4 people

Response Options

1 = None

2 = Some

3 = Much/Impossible

Variables Used:

ph110

DIStaste

Description: Respondent's self-rated sense of taste.

Response Options:

1 = Excellent

2 = Very Good

3 = Good

4 = Fair

5 = Poor or Registered Blind

Variables Used:

ph113

DISsmell

Description: Respondent's self-rated sense of smell.

Response Options:

1 = Excellent

2 = Very Good

3 = Good

4 = Fair

5 = Poor or Registered Blind

Variables Used:

ph112

Health

Physical and Mental Health

COGimmediatedrecall1

Description: Results of the first immediate recall question combining those answering from lists read out by the interviewer and a list read out from a computer.

Variables used:

ph117, ph119

COGimmediatedrecall2

Description: Results of the second immediate recall question combining those answering from lists read out by the interviewer and a list read out from a computer.

Variables used:

ph118, ph120

COGdelayedrecall

Description: Score for delayed recall test

Variables used:

ph712

COGprosmem1

Description: Result of prospective memory test 1 – record time

Response Options

0 = fail

1 = pass

Variables used:

Ph715

COGprosmem2

Description: Result of prospective memory test 2 – write initials

Response Options

0 = fail

1 = pass

Variables used:

ph419

COGmmse

Description: Total scoring of the Mini Mental State Examination.

Notes: The MMSE was only asked in the health assessment in Wave 1 of TILDA. It is included in the CAPI for Waves 2, 3, 4 and 5.

The total MMSE score was calculated by summing the item scores across all 11 tasks, taking the higher score of either the WORLD or Serial 7's (maximum 30 points).

For full details on the Mini-mental state examination, please refer to

<http://www.ncbi.nlm.nih.gov/pubmed/1202204>

A full comprehensive explanation of the MMSE scoring and tasks are included in the **TILDA Release Notes**.

Variables used

ph121, ph122, ph123, ph124, ph126, ph127, ph128, ph129, ph130, ph131, ph132, ph133, ph134, ph135, ph136, ph13, ph138, ph139, ph140,

COGanimal_naming

Description: Number of animals named in one minute

Note: Respondents are asked to name as many animals as they can in 1 minute

Variables used: ph125

MHcesd_capi_sf

Description: Indicator as to whether respondent has depressive symptoms.

Response options:

- 1 = Rarely or none of the time (less than 1 day)
- 2 = Some or a little of the time (1-2 days)
- 3 = Occasionally or a moderate amount of time (3-4 days)
- 4 = All of the time (5-7 days)

Scoring is as follows –

mh006, mh007, mh011, mh014, mh018, mh020,

The scale adds the value of the participant's response minus 1.

E.G. For question mh006, with the response "Occasionally or a moderate amount of time (3-4 days)", a 2 would be added to the scale result ($3 - 1 = 2$).

mh012, mh016

The scale adds 4 minus the value of the participant's response

E.G. For question mh012, with the response "Some or a little of the time (1-2 days)", a 2 would be added to the scale result ($4 - 2 = 2$)

Variables used: mh006, mh007, mh011, mh014, mh018, mh020

Further information:

O'Halloran, A. M., R. A. Kenny and B. L. King-Kallimanis (2014). "The latent factors of depression from the short forms of the CES-D are consistent, reliable and valid in community-living older adults." European Geriatric Medicine **5**(2): 97-102.

MHpennworry

Description: Overall result of the Penn State Worry scale.

Notes: SCQ only. Result is set to missing if any scale items have a missing response.

Response Options:

- 1 = Not at all typical
- 2 = In between
- 3 = Somewhat typical
- 4 = In between
- 5 = Very typical

Scoring:

The scale score is the result of all items added together. Each item is worth the value of their response option number.

Variables used:

SCQWORRY1 – SCQWORRY8

Further information:

Meyer, T. J., Miller, M. L., Metzger, R. L., & Borkovec, T. D. (1990). Development and validation of the Penn State Worry Questionnaire. *Behav Res Ther*, 28(6), 487-495.

MHucla_loneliness

Description: Overall result of the UCLA Loneliness scale.

Notes: SCQ only. Result is set to missing if any scale items have a missing response.

Response Options:

- 1 = Often
- 2 = Some of the time
- 3 = Hardly ever or never

Scoring:

Scoring is as follows –

SCQLoneLns1, 2, 3, 5:

The scale adds 3 minus the value of the participant's response.

E.G. For response "Often", a 2 would be added to the scale result ($3 - 1 = 2$).

SCQLoneLns4:

The scale adds value of the participant's response minus 1

E.G. For response "Some of the time", a 1 would be added to the scale result ($2 - 1 = 1$)

Variables used:

SCQLoneLns1 – SCQLoneLns5

Further information:

Russell, D. W. (1996). UCLA Loneliness Scale (Version 3): reliability, validity, and factor structure. *J Pers Assess*, 66(1), 20-40.

MHcapi_loneliness

Description: Frequency of loneliness taken from the CAPI questionnaire.

Notes: Don't know/Refused set to missing. Cloned variable of mh014.

1 = Rare or never

2 = Some of the time

3 = Moderate amount of the time

4 = All of the time

Variables used:

mh014

MHcidi_anxiety

Description: Indicates whether respondent fulfils the DSM criteria for general anxiety disorder based on an episode lasting 6 months or longer.

Notes: Uses the Composite International Diagnostic Interview – Short Form (CIDI).

For more information on the CIDI please refer to

<http://onlinelibrary.wiley.com/doi/10.1002/mpr.47/epdf>

DSM-IV criteria code: 300.02 - Generalized anxiety disorder

Criteria is met if participant has had –

Period of worry lasting at least 5 months (mh301, mh302, mh304m, mh304y, mh305m, mh305y, mh304, mh305)

Severe worry (mh306, mh307, mh308, mh310)

Difficult to control worry (mh309, mh311, mh312)

Additional miscellaneous anxiety symptoms (mh314 – mh320)

Response options:

0 = No

1 = Yes

Variables used:

mh301, mh302, mh304m, mh304y, mh305m, mh305y, mh304, mh305, mh306, mh307, mh308, mh309, mh310, mh311, mh312, mh314, mh315, mh316, mh317, mh318, mh319, mh320

MHcidi_depression

Description: Indicates whether respondent has had a major depressive episode in the past 12 months.

Notes: Uses the Composite International Diagnostic Interview – Short Form (CIDI).

For more information on the CIDI please refer to
<http://onlinelibrary.wiley.com/doi/10.1002/mpr.47/epdf>

DSM-IV criteria code: 300.4 Dysthymic disorder

There are two cases where participants are categorised as having a major depressive episode.

Participants who meet the criteria; has felt depressed or lost interest in most activities at least almost every day for at least most of the day in the past 12 months (mh101, mh102, mh103) are asked a number of additional depression symptoms (mh104, mh105, mh108, mh110, mh112, mh113). Those who meet the initial criteria and have 3 or more of the additional symptoms are coded as having had a major depressive episode.

Participants who meet the criteria; lost interest in most activities almost every day at least most of the day for two weeks or more in a row in the past 12 months (mh120, mh121, mh122) are asked a number of additional depression related symptoms (mh123, mh126, mh128, mh129, mh130, mh131). Those who meet the initial criteria and have 3 or more of the additional symptoms are coded as having had a major depressive episode.

Response options:

0 = No

1 = Yes

Variables used:

mh101, h102, mh103, mh120, mh121, mh122, mh104, mh105, mh108, mh110, mh112, mh113, mh123, mh126, mh128, mh129, mh130, mh131

MHcasp12_total

Description: Overall result including all domains of the CASP-19 scale.

Notes: SCQ only. CASP scale is set as missing if any question has a missing response.

Response Options:

1 = Often

2 = Not often

3 = Sometimes

4 = Never

Scoring is as follows –

SCQCASP1, 2, 4, 8 and 9 are reverse scored.

SCQCASP1, SCQCASP2, SCQCASP4, SCQCASP8, SCQCASP9:

The scale adds the value of the participant's response minus 1.

E.G. For question SCQCASP1, with the response "Not Often", a 1 would be added to the scale result ($2 - 1 = 1$).

SCQCASP3, SCQCASP7, SCQCASP10, SCQCASP11, SCQCASP13, SCQCASP17, SCQCASP18.

The scale adds 4 minus the value of the participant's response

E.G. For question SCQCASP3, with the response "Not often", a 2 would be added to the scale result ($4 - 2 = 2$)

Variables used:

SCQCASP1 – SCQCASP18

Further information:

Sexton, E., B. L. King-Kallimanis, R. M. Conroy and A. Hickey (2013). "Psychometric evaluation of the CASP-19 quality of life scale in an older Irish cohort." *Quality of Life Research* 22(9): 2549-2559. (<https://www.ncbi.nlm.nih.gov/pubmed/23504522>)

MHcasp12_control

Description: Result of the control domain of the CASP-12 scale.

Notes: SCQ only

Response Options:

1 = Often

2 = Not often

3 = Sometimes

4 = Never

The control domain includes the CASP1 – CASP4 items.

SCQCASP1, 2 and 4, are reverse scored.

Variables used:

SCQCASP1 – SCQCASP4

For more details on scoring, please refer to [MHcasp12_total](#)

MHcasp12_autonomy

Description: Result of the autonomy domain of the CASP-12 scale.

Notes: SCQ only

Response Options:

1 = Often

2 = Not often

3 = Sometimes

4 = Never

The control domain includes the CASP7 – CASP9 items.

SCQCASP8 and 9, are reverse scored.

Variables used:

SCQCASP7 – SCQCASP9

For more details on scoring, please refer to [MHcasp12_total](#)

MHcasp12_pleasure

Description: Result of the pleasure domain of the CASP-12 scale.

Notes: SCQ only

Response Options:

1 = Often

2 = Not often

3 = Sometimes

4 = Never

The control domain includes the CASP10 – CASP13 items.

Variables used:

SCQCASP10, SCQCASP11 and SCQCASP13

For more details on scoring, please refer to [MHcasp12_total](#)

MHcasp12_selfreal

Description: Result of the self-realisation domain of the CASP-12 scale.

Notes: SCQ only

Response Options:

1 = Often

2 = Not often

3 = Sometimes

4 = Never

The control domain includes the CASP17 – CASP18 items.

Variables used:

SCQCASP17 – SCQCASP18

For more details on scoring, please refer to [MHcasp12_total](#)

PSSscore

Description: Overall result including all domains of the Perceived Stress Scale (4items) 4-item. Scores range from 0 to 16, higher score indicates greater stress.

Notes: SCQ only. CASP scale is set as missing if any question has a missing response.

Response Options:

1 = Hardly ever

2 = Almost never

3 = Sometimes

4 = Fairly often

5 = Very often

Scoring is as follows –

SCQPPS2, 3 are reverse scored.

Variables used:

SCQPPS1, SCQPPS2, SCQPPS3, SCQPPS4

Further information:

Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24, 385-396.

CHRany_pain

Description: Indicates whether participant has any chronic pain

Note: includes proxy respondents

Response options:

0 = No pain

1 = Pain

Variables used:

Ph501

CHRpain_severity

Description: Indicates how severe participant's chronic pain is

Note: includes proxy respondents

Response options:

1 = Mild

2 = Moderate

3 = Severe

4 = Unknown

Variables used:

Ph502

CHRpainlimits

Description: Indicates whether the participants chronic pain limits their activities

Note: Excludes proxy respondents

Response options:

0 = No

1 = Yes

Variables used:

Ph504

MDmeds

Description: Number of reported medications (including supplements)

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDmeds4

Description: Number of reported medications (including supplements)

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDmeds_excl_supps

Description: Number of regular medications (excluding supplements)

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDmeds4_excl_supps

Description: Number of regular medications (excluding supplements)

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDsupplements

Description: Number of reported supplements

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDpolypharmacy

Description: Indicator variable for respondents who report more than five medications (including supplements)

Response options:

0 = No

1 = Yes

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDpoly_excl_supps

Description: Indicator variable for respondents who report more than five medications (excluding supplements)

Response options:

0 = No

1 = Yes

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDpoly_extreme

Description: Indicator variable for respondents who report more than ten medications (including supplements)

Response options:

0 = No

1 = Yes

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDpoly_extreme_ex_supps

Description: Indicator variable for respondents who report more than ten medications (excluding supplements)

Response options:

0 = No

1 = Yes

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDantidepressant

Description: Indicator variable for respondents who report at least one anti-depressant

Response options:

0 = No

1 = Yes

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDantihypertensives

Description: Indicator variable for respondents who report at least one anti-hypertensive (ATC C02 C07)

Response options:

0 = No

1 = Yes

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

MDantihypertensives2

Description: Indicator variable for respondents who report at least one anti-hypertensive (ATC C02 C03 C07 C08 C09)

Response options:

0 = No

1 = Yes

Variables used:

md001medcode_1 - md001medcode_20

md001medname_1 – md001medname_20

medcard

Description: Indicates whether respondent has a medical card.

Response options:

0 = No

1 = Yes

MDace_bbb

Description: Indicator of respondents who take a BBB (blood-brain barrier) crossing ACE inhibitor

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDace_nonbbb

Description: Indicator of respondents who take a non-BBB crossing ACE inhibitor

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDarb_bbb

Description: Indicator of respondents who take a BBB (blood-brain barrier) crossing ARB (Angiotensin II receptor blockers)

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDarb_nonbbb

Description: Indicator of respondents who take a non-BBB crossing ARB

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDarb_undbbb

Description: Indicator of respondents who take an undetermined ARB

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDccb_bbb

Description: Indicator of respondents who take a BBB crossing Calcium channel blocker

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDccb_nonbbb

Description: Indicator of respondents who take a non-BBB crossing Calcium channel blocker

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDantihypertensivesC08CEG

Description: Indicator of respondents who take an antihypertensive from ATC C08C, C08E, or C0

MDppi

Description: Taking any Proton pump inhibitor

Response options:

0 = No

1 = Yes

Variables used:

MDmeds, MDatc1-DMatc20

MDhypoglyc

Description: Taking any Blood glucose lowering drug, excl. insulins

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDinsulin

Description: Taking any Insulins and analogues

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MData1-DData20

MDglycosides

Description: Taking any Cardiac glycosides

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MData1-DData20

Mdantiarrhythmics

Description: Taking any Antiarrhythmics, class I and III

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MData1-DData20

MDvasodil

Description: Taking any Vasodilators used in cardiac diseases

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDantiadren

Description: Taking any Antiadrenergic agents, antihypertensives

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDdiur

Description: Taking any Diuretic use

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDlc_diur

Description: Taking any Low-ceiling diuretics

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDhc_diur

Description: Taking any High-ceiling diuretics

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDnons_betab

Description: Taking any Non-selective beta-blocker use

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDs_betab

Description: Taking any Selective beta blocker use

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDccb

Description: Taking any Calcium channel blockers

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDace

Description: Taking any ACE inhibitors

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDarb

Description: Taking any Angiotensin II Antagonist

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDstatin

Description: Taking any Statins

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDincont

Description: Taking any Drugs for urinary frequency and incontinence

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDalpha_bph

Description: Taking any Alpha-adrenoreceptor antagonist used in benign prostatic hypertrophy

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDnsaid

Description: Taking any NSAID

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDopioid

Description: Taking any Opioids

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDantiepileptic

Description: Taking any Antiepileptics

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDantipsych

Description: Taking any Antipsychotics

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDnons_mari

Description: Taking any Non-selective monoamine reuptake inhibitors

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDssri

Description: Taking any SSRIs

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDothertantidep

Description: Taking any Other Antidepressant

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDantihist

Description: Taking any Antihistamines

Notes: derived variables for the consortium for prediction model of drug-related falls

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDanticholinergic

Description: Taking any Anticholinergic medications

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDbenzodiazepine

Description: Taking any benzodiazepine

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDppi

Description: Taking any proton pump inhibitor

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

MDzed

Description: Taking any Z-drug hypnotic

Response options:

0 = No

1 = Yes

Variables used:

MDatc1-DMatc20

smokcurr

Description: Indicates whether respondent is a current smoker or not

Notes: Include all respondents who smoked in the past three months. Those who responded no in previous interviews are asked again and included if they now smoke.

Response options:

0 = No

1 = Yes

Variables used:

bh001, bh002, bh002a, bh002b

BEHcage

Description: CAGE alcohol scale score

Variables Used:

SCQCAGE1, SCQCAGE2, SCQCAGE3, SCQCAGE4

BEHcage2

Description: Indicator variable for respondents with an alcohol problem

Response options:

0 = No

1 = Yes

Variables Used:

SCQCAGE1, SCQCAGE2, SCQCAGE3, SCQCAGE4

ICD Codes

Incident cases of new diseases in Wave 3

TILDA Wave 3 uses the WHO International Statistical Classification of Diseases and Related Health Problems 10th Revision (ICD-10) to code incident cases of new conditions.

These variables apply only to new cases of conditions from Wave 2 to Wave 3. Any respondent with a condition is grouped into their relevant ICD code.

For more information on these and a full breakdown of all conditions included, please refer to <http://www.who.int/classifications/icd/en/>

ICD10_01

Description: Certain infectious and parasitic diseases

Response options:

0 = No

1 = Yes

Variables used:

N/A

ICD10_02

Description: Neoplasms

Response options:

0 = No

1 = Yes

Variables used:

ph301_05

ICD10_03

Description: Diseases of the blood

Response options:

0 = No

1 = Yes

Variables used:

N/A

ICD10_04

Description: Endocrine, nutritional and metabolic diseases

Response options:

0 = No

1 = Yes

Variables used:

ph201_05, ph201_08, ph301_15a

ICD10_05

Description: Mental and behaviour disorders

Response options:

0 = No

1 = Yes

Variables used:

ph301_07, ph301_08, ph301_08a, ph301_09, ph301_10, ph301_11

ICD10_06

Description: Diseases of the nervous system

Response options:

0 = No

1 = Yes

Variables used:

ph201_07, ph301_06

ICD10_07

Description: Diseases of the eye and adnexa

Response options:

0 = No

1 = Yes

Variables used:

ph105_01 - ph105_95

ICD10_08

Description: Diseases of the ear and mastoid process

Response options:

0 = No

1 = Yes

Variables used:

N/A

ICD10_09

Description: Diseases of the circulatory system

Response options:

0 = No

1 = Yes

Variables used:

ph201_01, ph201_02, ph201_03, ph201_04, ph201_06, ph201_09, ph201_10, ph201_11, ph201_95
ph301_13

ICD10_10

Description: Diseases of the respiratory system

Response options:

0 = No

1 = Yes

Variables used:

ph301_01, ph301_02

ICD10_11

Description: Diseases of the digestive system

Response options:

0 = No

1 = Yes

Variables used:

ph301_12, ph301_14

ICD10_12

Description: Diseases of the skin and subcutaneous tissue

Response options:

0 = No

1 = Yes

Variables used:

N/A

ICD10_13

Description: Diseases of the musculoskeletal system and connective tissue

Response options:

0 = No

1 = Yes

Variables used:

ph301_03, ph301_04

ICD10_14

Description: Diseases of the genitourinary system

Response options:

0 = No

1 = Yes

Variables used:

N/A

ICD10_15

Description: Conditions originating in the perinatal period

Response options:

0 = No

1 = Yes

Variables used:

N/A

ICD10_16

Description: Congenital malformations, deformations and chromosomal abnormalities

Response options:

0 = No

1 = Yes

Variables used:

N/A

IPAQ

Derived results from the IPAQ questionnaire

minswalking

Description: Number of minutes spent walking by respondent over past 7 days

Notes: Respondents are asked to record hours and minutes of time spent walking in the last 7 days. Hours are multiplied by 60 to generate overall minute count.

Includes any walking from place to place that lasted for 10 minutes or more.

Variables used:

bh106, bh106a

minsmoderate

Description: Number of minutes spent doing moderate physical activities by respondent in past 7 days

Notes: Respondents are asked to record hours and minutes of time spent doing moderate physical activities in the last 7 days. Hours are multiplied by 60 to generate overall minute count.

Moderate activity examples are carrying light loads, bicycling at a regular pace. Walking is not included

Variables used:

bh104, bh104a

minsvigorous

Description: Number of minutes spent doing vigorous physical activities by respondent in past 7 days

Notes: Respondents are asked to record hours and minutes of time spent doing vigorous physical activities in the last 7 days. Hours are multiplied by 60 to generate overall minute count.

Physical activity examples are heavy lifting, digging, aerobics or fast bicycling. Includes any vigorous activity that lasted for 10 minutes or more.

Variables used:

bh102, bh102a

walkingmet

Description: Met minutes spent walking by respondent over past 7 days

Notes: Amount of time walking (bh106, bh106a) is calculated by multiplying by the combined hours and minutes by the number of days the respondent has spent on these activities for at least 10 minutes at a time (bh105). This result is multiplied by 3.3.

Variables used:

bh105, bh106, bh106a

moderatemet

Description: Met minutes spent doing moderate physical activities by respondent in past 7 days

Notes: Amount of time spent on moderate activities (bh104, bh104a) is calculated by multiplying by the combined hours and minutes by the number of days the respondent has spent on these activities for at least 10 minutes at a time (bh103). This result is multiplied by 4.0.

Variables used:

bh103, bh104, bh104a

vigorousmet

Description: Met minutes spent doing vigorous physical activities by respondent in past 7 days

Notes: Amount of time spent on vigorous activities (bh102, bh102a) is calculated by multiplying by the combined hours and minutes by the number of days the respondent has spent on these activities for at least 10 minutes at a time (bh101). This result is multiplied by 8.0.

Variables used:

bh101, bh102, bh102a

FRipaquetminutes

Description: Total met (metabolic equivalent) minutes spent on vigorous activities, moderate activities and walking by the respondent in the past 7 days.

Notes: Respondent is asked how many hours and minutes per day they spend on these activities. Max minutes are set to 180.

Variables used:

bh101, bh102, bh102a, bh103, bh104, bh104a, bh105, bh106, bh106a

Further information:

http://www.institutferran.org/documentos/scoring_short_ipaq_april04.pdf

FRexercise3

Description: Categories of how much physical activity the respondent has engaged in.

Categories –

0 = Low

1 = Moderate

2 = High

Notes: Respondent is asked how many hours and minutes per day they spend on these activities. Max minutes are set to 180.

Scores are categorised as follows:

Moderate activity:

- Respondents who have spent 3 or more days doing 20 minutes of vigorous activity.
- Respondents who have spent 30 minutes or more walking and in moderate exercise for at least 5 days.
- Respondents who have 5 or more days of any activity totalling more than 600 MET minutes.

High Activity:

- Respondents who have spent 3 or more days on vigorous activity totalling 1500 or more MET minutes.
- Respondents who have spent 7 or more days on all activities totalling more than 3000 MET minutes.

Low activity:

- Respondents who have answered the questions but do not meet the criteria for either the moderate or high activity categories.

Variables used:

bh101, bh102, bh102a, bh103 bh104, bh104a, bh105, bh106, bh106a

Employment

Employment Status and Pensions

retired

Description: Indicates whether respondent is retired.

Response options:

0 = No

1 = Yes

Variables used:

we001

employ

Description: Indicates whether respondent is employed or self-employed.

Response options:

0 = No

1 = Yes

Variables used:

we001

pension_payIn

Description: Indicates whether respondent pays into any occupational, PRSA or private pensions.

Notes: Only includes respondents who are not retired.

Response options:

0 = No

1 = Yes

Variables used:

we001, wr002, we301, wr401

pension_receive

Description: Indicates whether respondent receives any pension payments – occupational, spousal, PSRA or private pension.

Response options:

0 = No

1 = Yes

Variables used:

si101, si201

Financial

Derived variables from financial questions

INCASSassets

Description: Value of gross assets

Variables used:

as101, as118, as103, as104, as106, as109, as121, as120, as112, as113, as116, hw105, hw113, hw109

INCASSnetassets

Description: Value of net assets

Variables used:

as101, as118, as103, as104, as106, as109, as121, as120, as112, as113, as116, hw105, hw113, hw109

INCASSassets5

Description: Value of gross assets in quintiles

Variables used:

as101, as118, as103, as104, as106, as109, as121, as120, as112, as113, as116, hw105, hw113, hw109

INCASSnetassets5

Description: Value of net assets in quintiles

Variables used:

as101, as118, as103, as104, as106, as109, as121, as120, as112, as113, as116, hw105, hw113, hw109

INCtotalhousehold_loop

Description: household income is based on a loop: the respondent is asked to estimate the income of each household member aged 16+, including himself/herself. Total household income is the sum of the income of each household member aged 16+. It is an estimation of NET income = after tax and deductions

Variables used:

si409_i, si412_i

INCtotalhousehold_loop

Description: household income in quintiles

Variables used:

si409_i, si412_i

Social Participation

SOCpaliveW5

Description: Indicates whether or not respondent has any living parents

Response Options

0 = None

1 = One Parent Alive

2 = Both

Variables used

tp001, tp007

SOCsocParticip

Description: Indicator of a whether or not respondents are in social groups

Response Options

0 = No

1 = Yes

Variables used

cn001

SOCChurch

Description: Whether or not a respondent attends religious services

Response Options

0 = No

1 = Yes

Variables used

dm021

SOCclubs

Description: Whether or not a respondent is involved in an organisation other than a Church

Response Option

0 = No

1 = Yes

Variables used

cn001

SOCrelFriends

Description: Count of close relatives or friends

Variables used

cn002, cn003, cn004

SOCcloseTies

Description: Whether respondent has at least 2 close friends or relatives

Response Options

0 = No

1 = Yes

Variables used

cn002, cn003, cn004

SOCsocnet

Description: Social Connectedness Score

Variables Used

cn001, cn002, cn003, cn004, cn006, dm021

SCR Section

Screening tests

SCRflu

Description: Indicator of whether respondent has ever had a flu vaccination.

Notes: Recode of ph701

0 = No

1 = Yes

Variables used:

ph701

SCRbreastlumps

Description: Indicator of whether respondent checks their breasts regularly for lumps.

Notes: Recode of ph708.

0 = No

1 = Yes

Variables used:

Ph708

SCRprostate

Description: Indicator of whether respondent has ever had prostate cancer.

Notes: Recode of ph711 and ph710. Respondents who have had either a prostate screen test for cancer or a PSA blood test are included.

0 = No

1 = Yes

Variables used:

ph710, ph711